

ALMA [C II]158 μ m DETECTION OF A REDSHIFT 7 LENSED GALAXY BEHIND RXJ1347.1-1145 *

MARUŠA BRADAČ¹, DIEGO GARCIA-APPADOO^{2,3}, KUANG-HAN HUANG¹, LIVIA VALLINI^{4,5}, EMILY QUINN FINNEY¹, AUSTIN HOAG¹, BRIAN C. LEMAUX¹, KASPER BORELLO SCHMIDT⁶, TOMMASO TREU^{7,x}, CHRIS CARILLI^{8,9}, MARK DIJKSTRA¹⁰, ANDREA FERRARA^{11,12}, ADRIANO FONTANA¹³, TUCKER JONES¹, RUSSELL RYAN¹⁴, JEFF WAGG¹⁵

Draft version October 11, 2016

ABSTRACT

We present the results of ALMA spectroscopic follow-up of a $z = 6.765$ Lyman- α emitting galaxy behind the cluster RXJ1347-1145. We report the detection of [C II]158 μ m line fully consistent with the Lyman- α redshift and with the peak of the optical emission. Given the magnification of $\mu = 5.0 \pm 0.3$ the intrinsic (corrected for lensing) luminosity of the [C II] line is $L_{[\text{C II}]} = 1.4^{+0.2}_{-0.3} \times 10^7 L_{\odot}$, which is ~ 5 times fainter than other detections of $z \sim 7$ galaxies. The result indicates that low $L_{[\text{C II}]}$ in $z \sim 7$ galaxies compared to the local counterparts are likely caused by their low metallicities and/or feedback. The small velocity off-set ($\Delta v = 20^{+140}_{-40}$ km s⁻¹) between the Lyman- α and [C II] line is unusual, and may be indicative of ionizing photons escaping.

Subject headings: galaxies: high-redshift — gravitational lensing: strong — galaxies: clusters: individual — dark ages, reionization, first stars

1. INTRODUCTION

The epoch of reionization, during which the Universe became transparent to UV radiation, is still poorly understood. Faint galaxies are likely responsible for this transformation, however this connection is far from confirmed (e.g. Robertson et al. 2015, Madau & Haardt 2015). Despite great progress in finding candidates with HST (e.g. Bouwens et al. 2015), beyond $z \gtrsim 7$, galaxies remain enshrouded in mystery, at least from a spectro-

scopic point of view. Spectroscopic confirmations remain extremely difficult, as the most prominent spectral feature in the optical/near IR wavelengths, the Lyman- α line, can be erased by neutral gas both in and surrounding galaxies.

An alternative way of measuring redshifts for early galaxies is to use sensitive radio/far-infra-red (FIR) telescopes to observe the [C II] 158 μ m line. It is among the strongest lines in star-forming galaxies at radio through FIR wavelengths (Carilli & Walter 2013) and it is therefore being actively pursued as a new way to measure redshifts at $z \gtrsim 6$. Furthermore, it allows us to study the connection of [C II] to Star Formation Rate (SFR), as well as the kinematics of distant galaxies (Knudsen et al. 2016c,b, Watson et al. 2015, Schaerer et al. 2015, Ota et al. 2014, Maiolino et al. 2015, Carilli & Walter 2013, Casey et al. 2014 and references therein). However, even with ALMA, detections at $z \sim 7$ remain sparse (see Pentericci et al. 2016, Maiolino et al. 2015, Knudsen et al. 2016c for current detections of [C II] and Inoue et al. 2016 for an [O III]88 μ m detection). In particular, despite deep observations, the extremely bright Lyman- α Emitter (LAE) Himiko was not detected in [C II] (Ouchi et al. 2013, Ota et al. 2014, Schaerer et al. 2015). From the theoretical side, the issue of non-detections has been studied e.g., by Vallini et al. (2015). The authors conclude that [C II] emission arises predominately from photodissociation regions (PDRs). Furthermore, the deficit in [C II]-SFR emission at $z \sim 7$ can be ascribed to either negative stellar feedback disrupting molecular clouds near the star forming regions and/or low gas metallicities.

In this paper we report on ALMA observations of RXJ1347:1216, a normal star forming galaxy with Lyman- α emission at $z_{\text{Ly}\alpha} = 6.7659^{+0.0030}_{-0.0005}$. It was first reported by Bradley et al. (2014) and Smit et al. (2014) as a photometrically selected $z \sim 7$ galaxy from the Hubble Space Telescope (*HST*) and *Spitzer* CLASH data (Postman et al. 2012). We have spectroscopically confirmed its redshift using Keck DEIMOS data (PI Bradač, Huang et al. 2016b) and *HST* grism data from GLASS

marusa@physics.ucdavis.edu

* These observations are based on the following ALMA data: ADS/JAO.ALMA#2015.1.0091.S. They are also associated with programs *Spitzer* #90009, 60034, 00083, 50610, 03550, 40593, and *HST* # GO10492, GO11591, GO12104, and GO13459. Furthermore based on multi-year KECK programs.

¹ Department of Physics, University of California, Davis, CA 95616, USA

² Joint ALMA Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile

⁴ European Southern Observatory, Alonso de Córdova 3107, Vitacura, Santiago, Chile

⁴ Dipartimento di Fisica e Astronomia, viale Berti Pichat 6, I-40127 Bologna, Italy

⁵ INAF, Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127 Bologna, Italy

⁶ Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany

⁷ Department of Physics and Astronomy, UCLA, Los Angeles, CA, 90095-1547, USA

⁸ National Radio Astronomy Observatory, P. O. Box 0, Socorro, NM 87801, USA

⁹ Cavendish Laboratory, 19 J. J. Thomson Ave., Cambridge CB3 0HE, UK

¹⁰ Institute of Theoretical Astrophysics, University of Oslo, Postboks 1029 Blindern, NO-0315 Oslo, Norway

¹¹ Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy

¹² Kavli IPMU (WPI), Todai Institutes for Advanced Study, the University of Tokyo, Japan

¹³ INAF - Osservatorio Astronomico di Roma Via Frascati 33 - 00040 Monte Porzio Catone, 00040 Rome, Italy

¹⁴ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

¹⁵ Square Kilometre Array Organisation, Lower Withington, Cheshire, UK

^x Packard Fellow

(PI Treu, Treu et al. 2015, Schmidt et al. 2016). In Huang et al. (2016b) we also improved the measurement of its stellar properties using deep *Spitzer* SURFSUP data (Bradač et al. 2014). Our magnification model shows that the massive foreground galaxy cluster RX J1347.1-1145 magnifies RXJ1347:1216 by a factor 5.0 ± 0.3 . Taking into account magnification, the galaxy’s intrinsic stellar mass is $M^* = 8.0_{-0.9}^{+6.5} \times 10^7 M_\odot$. The luminosity of RXJ1347:1216 is $0.2L^*$ (assuming characteristic magnitude at $z \sim 7$ of $M_{UV}^* = -20.87 \pm 0.26$ from Bouwens et al. 2015) making RXJ1347:1216 the first galaxy detected with ALMA at $z \sim 7$ having a luminosity characteristic of the majority of galaxies at $z \sim 7$.

The paper is structured as follows. In Section 2 we present the data and the reduction steps. In Section 3 we present the main science results as well as a detailed discussion of the findings. We summarize in Section 4. Throughout the paper we assume a Λ CDM concordance cosmology with $\Omega_m = 0.27$, $\Omega_\Lambda = 0.73$, and Hubble constant $H_0 = 73 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Coordinates are given for the epoch J2000.0, and magnitudes are in the AB system.

2. OBSERVATIONS AND DATA REDUCTION

We have observed RXJ1347:1216 with ALMA on 2016-06-21 as a part of the 2015.1.00091.S program (PI Bradač, priority A) in Band 6 with two executions. The total on-source integration time was 74 minutes with 38 antennas. The representative frequency was set to 244.92 GHz, and we requested angular resolution of $0''.3$ (the final beam size was $0''.59 \times 0''.41$). The phases were centered at the *HST* position of the source. The data were reduced with the Common Astronomy Software Application (CASA) and the images were produced using its subcomponent CLEAN. The continuum image was extracted using all the line-free channels of the four spectral windows.

3. RESULTS

We have detected [C II] emission of the LAE RXJ1347:1216 reported by Huang et al. (2016b) and Schmidt et al. (2016). The line is detected at 5σ and is spatially consistent (measured offset is $0''.1$) with the rest-frame UV emission (Fig. 1, Table 1). This offset is consistent with the standard ALMA astrometric uncertainty ($\sim 0''.1$; optical uncertainty is smaller). In addition, Dunlop et al. (2016) notes that the HST and ALMA astrometry of the HUDF field presented both a systematic shift of $0''.25$ and a random offset of up to $0''.5$. Similarly, Pentericci et al. (2016) notes $0''.1 - 0''.6$ random offsets measured from serendipitous detections in the field. Unfortunately we do not significantly detect any other sources in the field-of-view to perform relative astrometric calibration of the field.

The Lyman- α line redshift of this object is also in excellent agreement with the [C II] redshift (Fig. 2). The resulting velocity offset of Lyman- α compared to [C II] is only $\Delta v = 20_{-40}^{+140} \text{ km s}^{-1}$ (positive Δv indicates that Lyman- α is redshifted). The Lyman- α redshift was difficult to determine with a high accuracy given the proximity of the skyline, hence in Huang et al. (2016b) we report it only with two significant digits. The line is however significantly detected both in Keck/DEIMOS and in grism HST data from GLASS. We have since remeasured

TABLE 1
STELLAR POPULATION MODELING RESULTS FOR
RXJ 1347:1216 USING HST AND SPITZER
PHOTOMETRY FROM HUANG ET AL. (2016B),
SPECTROSCOPY OF LYMAN- α FROM KECK
DEMOS FROM HUANG ET AL. (2016B) AND
[C II] FROM ALMA OBSERVATIONS.

HST + Spitzer + Keck	
R.A.	13:47:36.207
Dec.	-11:45:15.16
$z_{\text{Ly}\alpha}$	6.7659 ± 0.0030
$EW_{\text{Ly}\alpha}$	$26 \pm 4 \text{ \AA}$
F160W ^a	$26.1 \pm 0.2 \text{ mag}$
μ_0	5.0 ± 0.3
$M_{\text{stellar}} \times f_\mu^b$	$8.0_{-0.9}^{+6.5} \times 10^7 M_\odot$
$SFR_{\text{SED}} \times f_\mu^b$	$8.5_{-1.0}^{+5.8} M_\odot \text{ yr}^{-1}$
$SFR_{\text{UV}} \times f_\mu^b$	$3.2 \pm 0.4 M_\odot \text{ yr}^{-1}$
Age	$\leq 13 \text{ Myr}$
$sSFR$	$105.1_{-20}^{+0.1} \text{ Gyr}^{-1}$
$E(B - V)_{\text{fit}}$	$0.10_{-0.01}^{+0.05}$
β	$-2.5_{-1.0}^{+0.7}$
ALMA	
R.A.	13:47:36.214
Dec.	-11:45:15.20
$\Delta_{\text{HST-ALMA}} [\text{arcsec}(\text{kpc})]$	0.1 (0.5)
$z_{[\text{CII}]}$	6.7655 ± 0.0005
S_{line}	$0.88 \pm 0.04 \text{ mJy}$
$S_{\text{line}} \Delta v$	$66 \pm 12 \text{ mJy km s}^{-1}$
FWHM	$75 \pm 25 \text{ km s}^{-1}$
$L_{[\text{CII}]} \times f_\mu^b$	$1.5_{-0.4}^{+0.2} 10^7 L_\odot$
Continuum	$< 15 \mu\text{Jy} (1\sigma)$
$L_{\text{FIR}} \times f_\mu^b$	$< 2 \times 10^{10} L_\odot (3\sigma)$
$SFR_{\text{FIR}} \times f_\mu^b$	$< 3 M_\odot \text{ yr}^{-1} (3\sigma)$

^a Lensed total magnitude (MAG_AUTO) in F160W.

^b The intrinsic properties calculated assuming $\mu = \mu_0$ from this table. To use a different magnification factor μ , simply use $f_\mu \equiv \mu/\mu_0$, i.e. dividing value given in this row by f_μ .




FIG. 1.— [C II] emission overlaid on the HST color composite RGB image (optical-F110W-F160W). The contours are spaced linearly between 0.4 – 0.9 mJy. A $1'' \times 1''$ zoom-in is shown in the inset, and the beam is given in bottom-left.

the redshift using our DEIMOS data (and improved on

absolute wavelength calibration as reported by Huang et al. 2016a). The reason for asymmetric errors on the measurement (Table 1) are the proximity of the skyline, low S/N of the line, and the asymmetric nature of the line. The errors are, however, small and we do not detect significant shift of the Lyman- α line.

This is unexpected, as for such a low luminosity galaxy the outflows are ubiquitous at lower- z . At $z \sim 3$ Erb et al. (2014) reports line red-shifts of $100 - 500 \text{ km s}^{-1}$ for low EW ($EW \lesssim 10$) LAEs and a strong anti-correlation between Δv and EW. The authors conclude that Δv is likely modulated both by galaxy continuum luminosity and by EW. In Huang et al. (2016b) we estimate the rest-frame EW to be $26 \pm 4 \text{ \AA}$. At $z \sim 7$, however, Stark et al. (2015) measure $\sim 60 \text{ km/s}$ for an offset between CIV and Lyman- α at $z = 7.045$ for a lower luminosity (lensed) object A1703-zd6. Pentericci et al. (2016) reports velocity shifts of $\sim 100 \text{ km/s}$ between [C II] and Lyman- α for their 3 most significant detections. For a higher luminosity galaxy Stark et al. (2016) reports an offset between CIII] and Lyman- α of $\sim 340 \text{ km/s}$. Given the lack of velocity offset of RXJ1347:1206 (and similar other low luminosity galaxies) it seems that at $z \sim 7$, Lyman- α is much closer to systemic velocity than is the case for low- z LAEs at similar luminosities.

This is important, since velocity offsets are crucial in interpreting the line visibility during the reionization epoch. The low offset is interpreted differently in the so called shell models vs. multi phase models of LAEs. A shell model requires low neutral gas column density or a high outflow velocity (Verhamme et al. 2015). In the case of a multi-phase model (Dijkstra et al. 2016), a low velocity off-set translates to a low covering factor of neutral gas, independent of its neutral column density and outflow velocity. In both models, however, the low velocity offset is a consequence of the presence of low HI-column density escape routes for Lyman- α photons, which may also allow for the escape of ionizing photons (see Verhamme et al. 2015, Dijkstra et al. 2016, Verhamme et al. 2016 for more detailed discussions). For a more general conclusion a larger sample is needed, but if future observations systematically show smaller velocity offsets, this would imply that the observed drop in LAE fraction between $z \sim 6$ and 7 (e.g. Schenker et al. 2012, 2014, Tilvi et al. 2014, Caruana et al. 2014, Pentericci et al. 2014, Schmidt et al. 2016) is more easily explained by changes in the IGM, than in the circumgalactic medium or galactic intrinsic properties (Dijkstra 2014, Mesinger et al. 2015, Choudhury et al. 2015).

The strength of the [C II] emission correlates with the SFR, however the [C II] luminosity is not a simple function of SFR, nor is there a simple dependence between [C II] luminosity and the total mass of the ISM. Recent simulations by Vallini et al. (2015), Pallottini et al. (2015), Vallini et al. (2016), Pallottini et al. (2016) indicate that [C II] emission at $z \sim 7$ arises predominately from PDRs, and that both feedback and low gas-phase metallicity appreciably lower the [C II] emission. This is shown in Fig. 3 where we plot $L_{[\text{C II}]}$ vs SFR. We include results from simulations by Vallini et al. (2015) as well as results for $z > 5$ galaxies from the literature. We also note, that we have estimated SFR using both rest-frame UV and optical data (we have also refit the photometric




FIG. 2.— Extracted spectrum (flux S_c as a function of frequency ν) of the [C II] emission. The red line denotes the best fit Gaussian (with parameter listed in Table 1 and the grey dashed line and region correspond to the Lyman- α redshift and uncertainty.

data from Huang et al. 2016b with a variety of metallicity templates; best fit result is shown in Fig. 4), hence it is more robust than SFRs determined using only UV photometry or based only on Lyman- α line. However, to compare results, we use SFR based on UV only in Fig. 3. The SED fitting results are compatible with models requiring some dust and consistent with the FIR measured SFR limits from the continuum non-detection (Table 1).

Due to gravitational lensing, we are able to measure [C II] luminosity in RXJ1347:1216 that is intrinsically ~ 5 times fainter than other such measurements at $z \sim 7$ to date (and similar luminosity to a $z \sim 6$ object in Knudsen et al. 2016a). The measurement is consistent with models requiring low metallicity $\lesssim 0.2 Z_\odot$, though feedback might play a role as well. As noted by Huang et al. (2016b), Smit et al. (2014), SED modelling for this galaxy (Fig. 4) requires strong nebular emission lines to explain its observed IRAC [3.6] – [4.5] colors. The object is best fit with young (10 Myr) stellar population, low metallicity ($0.2 Z_\odot$), and strong nebular emission lines ($EW > 100 \text{ \AA}$ for H β and O[III]; though note that all these parameters are highly degenerate); giving consistent picture with low [C II] luminosity compared to the local prediction. Recent detection of [O III]88 μm in a galaxy at $z = 7.2120$ with ALMA by Inoue et al. (2016) along with the upper limit on the [C II] is also in line with our results as it requires the ISM characterized by very low metallicity (Vallini et al. 2016, see also Cormier et al. 2015 for a study of local dwarf galaxies with high [O III]88 μm /[C II] ratios).

Following Pentericci et al. (2016), Wang et al. (2013) we can also estimate the dynamical mass of the galaxy. We use $M_{\text{dyn}} = 1.16 \times 10^5 V_{\text{cir}}^2 D$, where V_{cir} is the circular velocity in km/s, and D is the size in kpc, $V_{\text{cir}} = 0.75 FWHM_{[\text{C II}]} / \sin(i)$, and i is the disk inclination angle. Note that this approximation is very uncertain, as $z \sim 7$ galaxies might not have ordered motion. However, it was shown by Pallottini et al. (2016) that the [C II] mostly arises from an ordered, albeit small, disk. We estimate the size to be $\sim 0''.6$, and obtain $M_{\text{dyn}} \sin^2 i =$

$(1.0 \pm 0.5) \times 10^9 M_\odot$. For an intrinsically axisymmetric, infinitesimally thin disk, inclination is related to ellipticity via $\cos i = b/a$, where b/a is the axis ratio. With $b/a = 0.6$ measured from the line image we get $i = 50$ deg and we obtain $M_{\text{dyn}} = (2.0 \pm 0.9) \times 10^9 M_\odot$ (the error does not include systematic errors due to inclination). This value indicates that $f_{\text{gas}} \sim 1 - M^*/M_{\text{dyn}} = 96^{+2}_{-10} \%$ of the baryonic mass of this galaxy is in the molecular and atomic gas. The high value is consistent with predictions from the Kennicutt-Schmidt relation of SFR to gas mass (Kennicutt 1998), from which we obtain $M_{\text{gas}} = (3.5^{+2.1}_{-1.1}) \times 10^9 M_\odot$.




FIG. 3.— $L_{[\text{C II}]}$ vs SFR at $z > 5$. The red star is the $[\text{C II}]$ detection of the object RXJ 1347:1216. Lines represent the result from Vallini et al. (2015) obtained assuming a constant metallicity: black for $Z = Z_\odot$ (C1), orange for $Z = 0.2Z_\odot$ (C02), and blue for $Z = 0.05Z_\odot$ (C005). The magenta dashed line (P005) corresponds to a density-dependent metallicity with $Z = 0.05Z_\odot$. The 1σ scatter around the best-fit relations for dwarf and local starburst galaxies from De Looze et al. (2014) are plotted in dark gray and light gray, respectively. The green empty (filled) points represent upper limits (detections) of $[\text{C II}]$ in LAEs and LBGs at redshifts 5 – 7 as reported by Pentericci et al. (2016), Knudsen et al. (2016a), Inoue et al. (2016), Maiolino et al. (2015), Capak et al. (2015), Willott et al. (2015), Schaerer et al. (2015), González-López et al. (2014), Ota et al. (2014), Ouchi et al. (2013), Kanekar et al. (2013).

4. CONCLUSIONS

In this paper we report a $[\text{C II}]$ ALMA detection of a typical luminosity galaxy at $z = 6.765$. The $[\text{C II}]$ redshift agrees with Lyman- α redshift and the position agrees with the optical/UV *HST* counterpart of this object (within uncertainties). This indicates that the Lyman- α is at (or close to) resonant frequency, potentially requiring a lower IGM neutral fraction to explain the drop in the LAE fraction from $z \sim 7$ to $z \sim 6$ if the kinematics of the Ly α emitting region in RXJ1347:1216 are typical of $z \sim 7$ sub- L^* galaxies.




FIG. 4.— Updated SED fitting of HST+Spitzer photometry from Huang et al. (2016b). The galaxy is best fit model with a low metallicity template ($Z = 0.2Z_\odot$). We assume constant star formation history and fix the redshift to the spectroscopic redshift of the source.

The $[\text{C II}]$ luminosity is much lower than that expected from low-redshift low metallicity dwarf galaxies, yet it is consistent with predictions of simulations of $Z < 0.2Z_\odot$ galaxies at $z \sim 7$ by Vallini et al. (2015). This is the most likely explanation for why several searches for $[\text{C II}]$ emission at $z \sim 7$ have failed, as many have assumed local relation of $L_{[\text{C II}]}$ vs SFR derived for dwarf galaxies (De Looze et al. 2014). Due to gravitational lensing we reach lower intrinsic flux limits (factor of ~ 5) than similar observations of field galaxies, and as a result we are able to study a source that belongs to the bulk of the population at $z \sim 7$. As shown in this study, such lens magnified observations enable studies of the ISM in the sources responsible for reionization. Future higher resolution observations with ALMA will allow us to resolve (spectrally and spatially) galaxies at $z \sim 7$, and study in detail their kinematics and distribution of $[\text{C II}]$ at sub-kpc scales. Such studies will firmly establish ALMA as the premiere facility that will revolutionize the explorations of the earliest galaxies and our understanding of their place in the galaxy evolution and reionization puzzle.

We would like to thank Sergio Martin for help with the ALMA data. The work was carried out by MB while visiting Joint ALMA Observatory (JAO); MB acknowledges support for the visit by JAO through ALMA visitor programme. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2015.1.0091.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada), NSC and ASIAA (Taiwan), and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. Support was provided by NRAO via the NRAO Student Observing Support (SOS) Program. This material is based upon work supported by Associated Universities, Inc./National Radio Astronomy Observatory and the National Science Foundation Observations were also carried out using *Spitzer*

Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. Also based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555 and NNX08AD79G and W.M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation. Support for this work was provided by NASA

through an award issued by JPL/Caltech and through *HST*-AR-13235, *HST*-GO-13459, *HST*-GO-13177, *HST*-GO-10200, *HST*-GO-10863, and *HST*-GO-11099 from STScI. TT acknowledges support by the Packard Fellowship. AH acknowledges support by NASA Headquarters under the NASA Earth and Space Science Fellowship Program Grant ASTRO14F-0007. The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

Facilities: ALMA, Spitzer (IRAC), HST (ACS/WFC3), Keck/Deimos

REFERENCES

- Bouwens, R. J., Illingworth, G. D., Oesch, P. A., Trenti, M., Labbé, I., Bradley, L., Carollo, M., van Dokkum, P. G., Gonzalez, V., Holwerda, B., Franx, M., Spitler, L., Smit, R., & Magee, D. 2015, *ApJ*, 803, 34
- Bradač, M., Ryan, R., Casertano, S., Huang, K.-H., Lemaux, B. C., Schrabback, T., Gonzalez, A. H., Allen, S., Cain, B., Gladders, M., Hall, N., Hildebrandt, H., Hinz, J., von der Linden, A., Lubin, L., Treu, T., & Zaritsky, D. 2014, *ApJ*, 785, 108
- Bradley, L. D., Zitrin, A., Coe, D., Bouwens, R., Postman, M., Balestra, I., Grillo, C., Monna, A., Rosati, P., Seitz, S., Host, O., Lemze, D., Moustakas, J., Moustakas, L. A., Shu, X., Zheng, W., Broadhurst, T., Carrasco, M., Jouvel, S., Koekemoer, A., Medezinski, E., Meneghetti, M., Nonino, M., Smit, R., Umetsu, K., Bartelmann, M., Benítez, N., Donahue, M., Ford, H., Infante, L., Jimenez-Teja, Y., Kelson, D., Lahav, O., Maoz, D., Melchior, P., Merten, J., & Molino, A. 2014, *ApJ*, 792, 76
- Capak, P. L., Carilli, C., Jones, G., Casey, C. M., Riechers, D., Sheth, K., Carollo, C. M., Ilbert, O., Karim, A., Lefevre, O., Lilly, S., Scoville, N., Smolcic, V., & Yan, L. 2015, *Nature*, 522, 455
- Carilli, C. & Walter, F. 2013, *ArXiv:1301.0371*
- Caruana, J., Bunker, A. J., Wilkins, S. M., Stanway, E. R., Lorenzoni, S., Jarvis, M. J., & Ebert, H. 2014, *MNRAS*, 443, 2831
- Cassey, C. M., Narayanan, D., & Cooray, A. 2014, *Phys. Rep.*, 541, 45
- Choudhury, T. R., Puchwein, E., Haehnelt, M. G., & Bolton, J. S. 2015, *MNRAS*, 452, 261
- Cormier, D., Madden, S. C., Leboutteiller, V., Abel, N., Hony, S., Galliano, F., Rémy-Ruyer, A., Bigiel, F., Baes, M., Boselli, A., Chevance, M., Cooray, A., De Looze, I., Doublier, V., Galametz, M., Hughes, T., Karczewski, O. L., Lee, M.-Y., Lu, N., & Spinoglio, L. 2015, *A&A*, 578, A53
- De Looze, I., Cormier, D., Leboutteiller, V., Madden, S., Baes, M., Bendo, G. J., Boquien, M., Boselli, A., Clements, D. L., Cortese, L., Cooray, A., Galametz, M., Galliano, F., Graciá-Carpio, J., Isaak, K., Karczewski, O. L., Parkin, T. J., Pellegrini, E. W., Rémy-Ruyer, A., Spinoglio, L., Smith, M. W. L., & Sturm, E. 2014, *A&A*, 568, A62
- Dijkstra, M. 2014, *PASA*, 31, e040
- Dijkstra, M., Gronke, M., & Venkatesan, A. 2016, *ApJ*, 828, 71
- Dunlop, J. S., McLure, R. J., Biggs, A. D., Geach, J. E., Michalowski, M. J., Ivison, R. J., Rujopakarn, W., van Kampen, E., Kirkpatrick, A., Pope, A., Scott, D., Swinbank, A. M., Targett, T. A., Aretxaga, I., Austermann, J. E., Best, P. N., Bruce, V. A., Chapin, E. L., Charlot, S., Cirasuolo, M., Coppin, K. E. K., Ellis, R. S., Finkelstein, S. L., Hayward, C. C., Hughes, D. H., Ibar, E., Khochfar, S., Koproński, M. P., Narayanan, D., Papovich, C., Peacock, J. A., Robertson, B., Vernstrom, T., van der Werf, P. P., Wilson, G. W., & Yun, M. 2016, *ArXiv:1606.00227*
- Erb, D. K., Steidel, C. C., Trainor, R. F., Bogosavljević, M., Shapley, A. E., Nestor, D. B., Kulas, K. R., Law, D. R., Strom, A. L., Rudie, G. C., Reddy, N. A., Pettini, M., Konidakis, N. P., Mace, G., Matthews, K., & McLean, I. S. 2014, *ApJ*, 795, 33
- González-López, J., Riechers, D. A., Decarli, R., Walter, F., Vallini, L., Neri, R., Bertoldi, F., Bolatto, A. D., Carilli, C. L., Cox, P., da Cunha, E., Ferrara, A., Gallerani, S., & Infante, L. 2014, *ApJ*, 784, 99
- Huang, K. H., Bradač, M., Allen, S., Cain, B., Casertano, S., Gladders, M., Gonzalez, A. H., Hoag, A., Hildebrandt, H., Hinz, J., Lemaux, B. C., von der Linden, A., Lubin, L., Ryan, R., Schrabback, T., Treu, T., & Zaritsky, D. 2016a, in preparation
- Huang, K.-H., Bradač, M., Lemaux, B. C., Ryan, Jr., R. E., Hoag, A., Castellano, M., Amorín, R., Fontana, A., Brammer, G. B., Cain, B., Lubin, L. M., Merlin, E., Schmidt, K. B., Schrabback, T., Treu, T., Gonzalez, A. H., von der Linden, A., & Knight, R. I. 2016b, *ApJ*, 817, 11
- Inoue, A. K., Tamura, Y., Matsuo, H., Mawatari, K., Shimizu, I., Shibuya, T., Ota, K., Yoshida, N., Zackrisson, E., Kashikawa, N., Kohno, K., Umehata, H., Hatsukade, B., Iye, M., Matsuda, Y., Okamoto, T., & Yamaguchi, Y. 2016, *Science*, 352, 1559
- Kanekar, N., Wagg, J., Ram Chary, R., & Carilli, C. L. 2013, *ApJ*, 771, L20
- Kennicutt, Jr., R. C. 1998, *ApJ*, 498, 541
- Knudsen, K. K., Richard, J., Kneib, J.-P., Jauzac, M., Clément, B., Drouart, G., Egami, E., & Lindroos, L. 2016a, *MNRAS*, 462, L6
- Knudsen, K. K., Richard, J., Kneib, J.-P., Jauzac, M., Clément, B., Drouart, G., Egami, E., & Lindroos, L. 2016b, *ArXiv:1603.02277*
- Knudsen, K. K., Watson, D., Frayer, D., Christensen, L., Gallazzi, A., Michalowski, M. J., Richard, J., & Zavala, J. 2016c, *ArXiv:1603.03222*
- Madau, P. & Haardt, F. 2015, *ApJ*, 813, L8
- Maiolino, R., Carniani, S., Fontana, A., Vallini, L., Pentericci, L., Ferrara, A., Vanzella, E., Grazian, A., Gallerani, S., Castellano, M., Cristiani, S., Brammer, G., Santini, P., Wagg, J., & Williams, R. 2015, *ArXiv:1502.06634*
- Mesinger, A., Aykutalp, A., Vanzella, E., Pentericci, L., Ferrara, A., & Dijkstra, M. 2015, *MNRAS*, 446, 566
- Ota, K., Walter, F., Ohta, K., Hatsukade, B., Carilli, C. L., da Cunha, E., González-López, J., Decarli, R., Hodge, J. A., Nagai, H., Egami, E., Jiang, L., Iye, M., Kashikawa, N., Riechers, D. A., Bertoldi, F., Cox, P., Neri, R., & Weiss, A. 2014, *ApJ*, 792, 34
- Ouchi, M., Ellis, R., Ono, Y., Nakanishi, K., Kohno, K., Momose, R., Kurono, Y., Ashby, M. L. N., Shimasaku, K., Willner, S. P., Fazio, G. G., Tamura, Y., & Iono, D. 2013, *ApJ*, 778, 102
- Pallottini, A., Ferrara, A., Gallerani, S., Vallini, L., Maiolino, R., & Salvadori, S. 2016, *ArXiv:1609.01719*
- Pallottini, A., Gallerani, S., Ferrara, A., Yue, B., Vallini, L., Maiolino, R., & Feruglio, C. 2015, *MNRAS*, 453, 1898
- Pentericci, L., Carniani, S., Castellano, M., Fontana, A., Maiolino, R., Guaita, L., Vanzella, E., Grazian, A., Santini, P., Yan, H., Cristiani, S., Conselice, C., Giallisco, M., Hathi, N., & Koekemoer, A. 2016, *ArXiv:1608.08837*
- Pentericci, L., Vanzella, E., Fontana, A., Castellano, M., Treu, T., Mesinger, A., Dijkstra, M., Grazian, A., Bradač, M., Conselice, C., Cristiani, S., Dunlop, J., Galametz, A., Giallisco, M., Giallongo, E., Koekemoer, A., McLure, R., Maiolino, R., Paris, D., & Santini, P. 2014, *ApJ*, 793, 113
- Postman, M., Coe, D., Benítez, N., Bradley, L., Broadhurst, T., Donahue, M., Ford, H., Graur, O., Graves, G., Jouvel, S., Koekemoer, A., Lemze, D., Medezinski, E., Molino, A., Moustakas, L., Ogaz, S., Riess, A., Rodney, S., Rosati, P., Umetsu, K., Zheng, W., Zitrin, A., Bartelmann, M., Bouwens, R., Czakon, N., Golwala, S., Host, O., Infante, L., Jha, S., Jimenez-Teja, Y., Kelson, D., Lahav, O., Lazkoz, R., Maoz, D., McCully, C., Melchior, P., Meneghetti, M., Merten, J., Moustakas, J., Nonino, M., Patel, B., Regös, E., Sayers, J., Seitz, S., & Van der Wel, A. 2012, *ApJS*, 199, 25
- Robertson, B. E., Ellis, R. S., Furlanetto, S. R., & Dunlop, J. S. 2015, *ApJ*, 802, L19
- Schaerer, D., Boone, F., Zamojski, M., Staguhn, J., Dessauges-Zavadsky, M., Finkelstein, S., & Combes, F. 2015, *A&A*, 574, A19

- Schenker, M. A., Ellis, R. S., Konidaris, N. P., & Stark, D. P. 2014, ArXiv:1404.4632
- Schenker, M. A., Stark, D. P., Ellis, R. S., Robertson, B. E., Dunlop, J. S., McLure, R. J., Kneib, J.-P., & Richard, J. 2012, ApJ, 744, 179
- Schmidt, K. B., Treu, T., Bradač, M., Vulcani, B., Huang, K.-H., Hoag, A., Maseda, M., Guaita, L., Pentericci, L., Brammer, G. B., Dijkstra, M., Dressler, A., Fontana, A., Henry, A. L., Jones, T. A., Mason, C., Trenti, M., & Wang, X. 2016, ApJ, 818, 38
- Smit, R., Bouwens, R. J., Labbé, I., Zheng, W., Bradley, L., Donahue, M., Lemze, D., Moustakas, J., Umetsu, K., Zitrin, A., Coe, D., Postman, M., Gonzalez, V., Bartelmann, M., Benítez, N., Broadhurst, T., Ford, H., Grillo, C., Infante, L., Jimenez-Teja, Y., Jouvel, S., Kelson, D. D., Lahav, O., Maoz, D., Medezinski, E., Melchior, P., Meneghetti, M., Merten, J., Molino, A., Moustakas, L. A., Nonino, M., Rosati, P., & Seitz, S. 2014, ApJ, 784, 58
- Stark, D. P., Ellis, R. S., Charlot, S., Chevallard, J., Tang, M., Belli, S., Zitrin, A., Mainali, R., Gutkin, J., Vidal-Garcia, A., Bouwens, R., & Oesch, P. 2016, ArXiv:1606.01304
- Stark, D. P., Walth, G., Charlot, S., Clément, B., Feltre, A., Gutkin, J., Richard, J., Mainali, R., Robertson, B., Siana, B., Tang, M., & Schenker, M. 2015, MNRAS, 454, 1393
- Tilvi, V., Papovich, C., Finkelstein, S. L., Long, J., Song, M., Dickinson, M., Ferguson, H. C., Koekemoer, A. M., Giavalisco, M., & Mobasher, B. 2014, ApJ, 794, 5
- Treu, T., Schmidt, K. B., Brammer, G. B., Vulcani, B., Wang, X., Bradač, M., Dijkstra, M., Dressler, A., Fontana, A., Gavazzi, R., Henry, A. L., Hoag, A., Huang, K.-H., Jones, T. A., Kelly, P. L., Malkan, M. A., Mason, C., Pentericci, L., Poggianti, B., Stiavelli, M., Trenti, M., & von der Linden, A. 2015, ApJ, 812, 114
- Vallini, L., Ferrara, A., Pallottini, A., & Gallerani, S. 2016, ArXiv:1606.08464
- Vallini, L., Gallerani, S., Ferrara, A., Pallottini, A., & Yue, B. 2015, ApJ, 813, 36
- Verhamme, A., Orlitová, I., Schaerer, D., & Hayes, M. 2015, A&A, 578, A7
- Verhamme, A., Orlitova, I., Schaerer, D., Izotov, Y., Wörseck, G., Thuan, T. X., & Guseva, N. 2016, ArXiv:1609.03477
- Wang, R., Wagg, J., Carilli, C. L., Walter, F., Lentati, L., Fan, X., Riechers, D. A., Bertoldi, F., Narayanan, D., Strauss, M. A., Cox, P., Omont, A., Menten, K. M., Knudsen, K. K., Neri, R., & Jiang, L. 2013, ApJ, 773, 44
- Watson, D., Christensen, L., Knudsen, K. K., Richard, J., Gallazzi, A., & Michałowski, M. J. 2015, Nature, 519, 327
- Willott, C. J., Carilli, C. L., Wagg, J., & Wang, R. 2015, ApJ, 807, 180